# Taking Itulazax® for severe birch pollen allergy

Adult Allergy service	Produced:	December 2023
	Updated:	November 2024
	Review:	December 2026
Information for Patients	Leaflet number: 1455 Version: 1.1	

You should read this leaflet together with the manufacturer's leaflet for Itulazax®.

You will get a copy of the manufacturer's guide with your supply of Itulazax®.Copies are also available from <u>www.medicines.org.uk/emc/</u>

#### What is Itulazax<sup>®</sup> and how is it taken?

Itulazax® is used to treat birch tree pollen allergy in adults who are living with moderate to severe symptoms– despite taking other symptom relieving medications, such as antihistamine tablets, nose sprays and eye drops.

- You will be prescribed the same dose. You take it every day for 3 years.
- At your 1st appointment you will be given a prescription for 4 week supply of Itulazax® tablets.
- You should take your antihistamine tablet (if not already taking) on the day you start Itulazax® and for 2 weeks after starting.
- Please take your **tablet every single day** (at the same time) even when you are well. Please read the manufacturer's leaflet for instructions on how to take Itulazax®.
- You should **keep on taking** all your medicines like antihistamines, nasal spray and eye drops as normal after starting Itulazax®.
- If you forget your Itulazax® tablet, take it later that day. If you forget that day than continue with the one a day routine on the next day. Do not take a double dose in 1 day.
- In most cases, people start to feel better after taking Itulazax® once a day in the tree pollen season.

#### Health information and support is available at www.nhs.uk or call 111 for non-emergency medical advice

Visit www.leicestershospitals.nhs.uk for maps and information about visiting Leicester's Hospitals To give feedback about this information sheet, contact InformationForPatients@uhl-tr.nhs.uk



## Who should not take Itulazax?

Do not take Itulazax if:

- You have uncontrolled asthma
- You are pregnant or breast feeding

Itulazax may not be suitable if:

- You are allergic to fish or do not eat fish
- You have a condition that affects your immune system
- You have cancer
- You have previously had a severe allergic reaction to injection immunotherapy

### What should I do if I have a reaction to Itulazax®?

If you experience any potentially severe side effects, you should seek emergency treatment by **DIALLING 999. DO NOT TAKE ANYMORE ITULAZAX.** Examples of severe side effects include:

- Difficulty in swallowing
- Difficulty in breathing
- Change in your voice (hoarse voice)
- Rapid swelling of face, mouth, throat or skin
- Severe dizziness

Please check the manufacturer's leaflet for more information on possible side effects.

You should expect some mild to moderate local allergic reactions in the first few days. These usually start 5 to 10 minutes after taking the Itulazax and only last a few minute or hours. They should lessen over the first few months as you continue treatment.

If any side effects are worrying you or causing you problems please contact us. Do not take any further doses until you have spoken to the allergy nurses.

Examples of very common and common side effects are:

- Itching of the ears and mouth
- Sneezing
- Swelling or itching under the tongue (where the Itulazax® tablet is put)
- Runny nose

#### Other reasons to stop Itulazax®

- You should stop Itulazax for a limited time if you have problems with your mouth, such as soreness, bleeding, infections or have had dental treatment. Restart after your mouth is totally healed.
- You have an infection which makes you feel unwell or gives you a high temperature
- You are having any vaccinations do not take your Itulazax on the same day as your vaccination.
- Your consultant may stop your this treatment if you do not take it as directed or if you do not attend your follow-up appointments.

If you need to stop Itulazax®:

- For 1 week or less, you can restart at home but please call the nursing team to tell us why you had stopped and for how many days.
- For more than 1 week, you must call and speak to the nursing team before restarting your treatment. You may have to restart treatment back in the hospital.
- For more than 4 weeks it is not safe to restart at home. Please call the allergy nurses to tell us why it has been stopped and for how many days. If appropriate, it can be re started in clinic but only after being discussed with your doctor.

#### How do I order more prescriptions of Itulazax®?

A Shared Care Agreement (SCA) is an agreement between you, your GP and your hospital consultant. The agreement means that you can keep on taking the medication your doctor has prescribed instead of going to the hospital to pick it up.

Once your GP has agreed to the SCA, the GP will be able to prescribe the Itulazax® for you. Remember to ask for new prescriptions from your GP so that there are no gaps in your treatment.

If your GP has **not agreed** to a SCA you will need to collect your medicine from the hospital pharmacy (Trustmed Pharmacy). Please call the allergy nurses at the hospital to order when you have 2 weeks supply left. The nurses are not able to order more treatment until they have spoken to you. Trustmed Pharmacy will call you when your medication is ready for you to pick up.

#### When will I have follow-up appointments?

You will be reviewed by our allergy consultants every year.

You will also have follow-up appointment with our allergy nurses once a year.

At these appointments, you will be

- checked to see how you are doing with the treatment
- advised if the treatment needs to carry on or not



• Your consultant may stop this treatment if you do not take it as directed, or your symptoms do not improve, or if you do not attend your follow-up appointments.

If you get better with Itulazax and you take it regularly, you will carry on being on this treatment for 3 years.

#### **Contact details:**

Allergy nursing service contact details for patients started at and under the care of Glenfield Hospital 0116 258 3557

Allergy nursing service contact details for patients started at and under the care of Leicester General Hospital 0116 258 6702 or 0116 258 6711

Monday to Friday, 8am to 4.00pm (message can be left on our answerphone when we are not available and we will call you back). Please speak to nursing team to order your medication.

# **Further information**

LEICESTER'S

Please find more information on allergies on Allergy UK website: www.allergyuk.org

اگر آپ کو یہ معلومات کسـی اور زبان میں درکار ہیں، تو براہِ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔ علی هذه المعلومات بلغةٍ أُخری، الرجاء الاتصال علی رقم الهاتف الذي يظهر في الأسـفل જો તમને અન્ય ભાષામાં આ માફિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો

ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ `ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ। Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email equality@uhl-tr.nhs.uk

Leicester's Hospitals is a research active trust so you may find research happening on your ward or in your clinic. To find out about the benefits of research and become involved yourself, speak to your clinician or nurse, call 0116 258 8351 or visit www.leicestersresearch.nhs.uk/ patient-and-public-involvement